



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Virus structure and structure-based antivirals

Zlatka Plavec^{1,2}, Ina Pöhner³, Antti Poso^{3,4} and Sarah J Butcher^{1,2}

Structure-based antiviral developments in the past two years have been dominated by the structure determination and inhibition of SARS-CoV-2 proteins and new lead molecules for picornaviruses. The SARS-CoV-2 spike protein has been targeted successfully with antibodies, nanobodies, and receptor protein mimics effectively blocking receptor binding or fusion. The two most promising non-structural proteins sharing strong structural and functional conservation across virus families are the main protease and the RNA-dependent RNA polymerase, for which design and reuse of broad range inhibitors already approved for use has been an attractive avenue. For picornaviruses, the increasing recognition of the transient expansion of the capsid as a critical transition towards RNA release has been targeted through a newly identified, apparently widely conserved, druggable, interprotomer pocket preventing viral entry. We summarize some of the key papers in these areas and ponder the practical uses and contributions of molecular modeling alongside empirical structure determination.

Addresses

¹ Faculty of Biological and Environmental Sciences, Molecular and Integrative Bioscience Research Programme, University of Helsinki, Helsinki, Finland

² Helsinki Institute of Life Sciences-Institute of Biotechnology, University of Helsinki, Helsinki, Finland

³ School of Pharmacy, University of Eastern Finland, Kuopio, Finland

⁴ University Hospital Tübingen, Department of Internal Medicine VII, Tübingen, Germany

Corresponding authors:

Plavec, Zlatka (zlatka.plavec@helsinki.fi), Butcher, Sarah J (sarah.butcher@helsinki.fi)

Current Opinion in Virology 2021, 51:16–24

This review comes from a themed issue on **Virus structure and expression**

Edited by **José R Castón** and **Adam Zlotnick**

For complete overview about the section, refer "[Virus structure and expression \(2022\)](#)"

Available online 24th September 2021

<https://doi.org/10.1016/j.coviro.2021.09.005>

1879-6257/© 2021 Published by Elsevier B.V.

Introduction

Since the beginning of the Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) pandemic in

2019, there has been a race between antiviral drug design and vaccine development to combat the disease. Although vaccines won the race, lowering the risk of severe infection, there is still a need for antiviral drugs, driving research for SARS-CoV-2, and other significant viral pathogens. The literature in the past two years has been dominated by studies utilizing computational methods to identify SARS-CoV-2 antivirals [1–15]. In this Opinion we focus on SARS-CoV-2 and picornaviruses, and antiviral leads where structural data have contributed significantly to the design or interpretation of the antiviral mechanism and led to insightful information for future structure-based drug design.

SARS-CoV-2

SARS-CoV-2 belongs to the *Coronaviridae* family, a family of pleiomorphic ss(+)RNA viruses. Its genome is associated with the N protein forming the nucleocapsid, coated with a spike glycoprotein-adorned membrane [16]. The most promising drug targets against SARS-CoV-2 have been the spike glycoprotein (S) and the non-structural components: the main protease (M^{Pro}), the papain like protease (PL^{Pro}), and the RNA-dependent RNA polymerase (RdRp) [7,17–23]. For the non-structural, highly conserved proteins, drug repurposing and broad range inhibition was an attractive approach [3,11,18–22,24–28,29*,30**,31–45]. S, as a surface protein is antigenic and is prone to mutations [17]. A complementary strategy of inhibiting host proteins required in infection is exemplified by the use of soluble peptides mimicking the angiotensin-converting enzyme (ACE2) receptor binding domain (RBD) [46,47].

Picornaviruses

The noticeable structural and functional conservation between the proteases and polymerases of SARS-CoV-2 and those of picornaviruses has led to research focusing on broad-range inhibitors targeting both families. Picornaviruses are a family of small, 30 nm icosahedrally symmetric viruses with (+)ssRNA genome. The capsids consist of three β-barrel viral proteins (VP1, VP2, and VP3) and one elongated internal protein, VP4. Many of the viruses in the family have a conserved hydrophobic pocket in VP1 that can contain a lipid factor [48,49]. Several picornavirus inhibitors target this conserved pocket, preventing genome release, but none of the inhibitors are currently in clinical use [50]. We will describe advances targeting a novel interprotomer capsid pocket instead.


Drug-repurposing and computational structure-based drug design

Drug-repurposing has rapidly advanced SARS-CoV-2 antiviral approaches avoiding lengthy clinical drug approval processes, building on work done to inhibit SARS-CoV, MERS-CoV, hepatitis C and so on. Amongst the drugs which have shown the most promising results and have gone on to clinical trials are the hepatitis C antivirals, telaprevir, narnavir and boceprevir inhibiting M^{pro} and remdesivir, ribavirin, suramin and favipiravir riboside triphosphate (RTP) against the RdRp [31–34,44,51]. SARS-CoV-2 M^{pro} was a very promising target at the start of the pandemic and its potential and the advances made have been extensively reviewed: Mengist *et al.* reviewed their toxicology and the structural basis for their inhibition while some other reviews have focused on their potential for clinical use and future prospects [18–22,39,40]. Like many other non-structural proteins, M^{pro} is highly conserved in structure and substrate specificity across the *Coronaviridae* family as well as some other virus families, like the *Picornaviridae*. Both M^{pro} and the picornaviral $3C^{pro}$ have chymotrypsin-like folds and cleave the viral polyprotein into functional proteins, which in turn are involved in viral replication. SARS-CoV-2 M^{pro} is a homodimer with a Cys-His catalytic dyad between two domains (Figure 1a). M^{pro} compared to $3C^{pro}$ has an additional α -helical domain connected by a loop to the catalytic site. Its substrate-binding pocket subsites and the corresponding inhibitor positions are shown in Figure 1c. Unlike SARS-CoV-2 M^{pro} , $3C^{pro}$ is a monomer and has a Cys-His-Glu/Asp catalytic triad (Figure 1d). Both proteases require Gln as the P1 residue in the substrate and only have space for a small amino-acid residue at P1' (Figure 1) [30^{**},35,36]. For this reason, inhibitors targeting SARS-CoV M^{pro} and picornavirus $3C^{pro}$ were used as templates for design of SARS-CoV-2 M^{pro} [24,26,30^{**},35–38,41,52,53].

Over 1000 M^{pro} structures have now been deposited in the PDB giving an unprecedented opportunity for detailed data mining. Although M^{pro} is catalytically active as a homodimer, almost all modeling studies have been based on the monomeric structure. Weng *et al.* (2021), however, utilized and validated the effect of dimers for inhibitor binding on the molecular level [54]. The initial excitement over repurposing of protease inhibitors has so far not led to effective clinical use and this may well be partially due to the little knowledge on how dimerization affects the inhibitor binding region.

The RdRp has also been explored as a target for drug repurposing [18,20,27,29^{*},43,45]. It is a vital enzyme transcribing RNA from its complementary template. Structurally, it resembles other right-hand polymerases consisting of 3 conserved domains termed the palm, fingers and thumb (Figure 2). The structural similarity

Figure 1




Current Opinion in Virology

Structure of SARS-CoV-2 M^{pro} and coxsackievirus B3 $3C^{pro}$ bound with inhibitors.

(a) Surface representation of SARS-CoV-2 M^{pro} (PDB: 6LU7) with monomers in blue and magenta. Catalytic dyad is marked in yellow and inhibitor in cyan. (b) Close up of the active site residues and interactions with the inhibitor N3 [30^{**}]. (c) Close up of the active site and the inhibitor N3 showing active site subsites with yellow circles and inhibitor positions marked P1'-P5 [30^{**}]. (d) Surface representation of coxsackievirus B3 $3C^{pro}$ in magenta with inhibitor 11a in cyan and catalytic triad in yellow (PDB: 3ZYD) [24]. (e) Close up of the active site residues and interactions with inhibitor 11a. (f) Close up of the active sites and the inhibitor 11a showing active site subsites with yellow circles and inhibitor positions marked P1'-P3 [24].

Figure created in UCSF Chimera [87]. (M^{pro} = main protease, $3C^{pro}$ = 3C protease).

of the RdRp across families of RNA viruses was recently used by Mönttinen *et al.* to create a phylogenetic tree. SARS-CoV-2 RdRp clusters with several flaviviruses and hepaciviruses, adding weight to repurposing the previously mentioned inhibitors remdesivir, suramin and ribavirin, all developed as hepatitis C antivirals [25]. Structural characterization of the SARS-CoV-2 RdRp started with the description of the complex with accessory factors nsp7 and nsp8 [27,28]. Complexes with the RNA template and finally with the nucleotide-analog remdesivir (Figure 2) progressively improved the understanding of RdRp activity and the mechanism of nucleotide-based


Figure 2

SARS-CoV-2 RdRp with nsp7, nsp8, RNA template and remdesivir. (a) Surface and (b) ribbon view of RdRp structure with color-coded domains labelled. (c) Close up view of remdesivir and its location in the RdRp. Pyrophosphate (PP) is marked in orange and is a byproduct of the remdesivir-inhibited reaction [34]. Figures created in UCSF Chimera [87] from PDB:7BV2. Nidovirus RdRp-associated nucleotidyl transferase domain (NiRAN).

inhibitors [27,34]. A cryo-EM structure with suramin further indicates the potential for distinct chemical entities against SARS-CoV-2 RdRp [44]. The structural conservation between SARS-CoV-2 and flavivirus RdRp indicates a potential alternative option for drug discovery: two inhibitors targeting the flexible RNA tunnel have been described for Dengue virus RdRp. The interacting residues are largely conserved across flaviviruses and may allow the design of broad-spectrum antivirals along with those targeting the active site [55,56]. While the lack of flavivirus RdRp structures in their catalytically competent state represents a challenge for structure-based drug discovery approaches, a recent structure of TBEV RdRp helped to identify the differences in catalytic sites between tick-borne and mosquito-borne flaviviruses. These differences were shown to be vital in host recognition, expanding the known RdRp roles in the infection cycle [45,57].

Structural proteins as targets

Viral structural proteins are the first to interact with the host whether by interacting with other proteins such as receptors or antibodies, or the host environment in the form of pH, ions or lipids. As such, they are prone to mutations to evade the host immune system. In SARS-CoV-2 infection, S protein attaches to the host receptor,

Figure 3

SARS-CoV-2 spike glycoprotein conformational states, glycosylation and binding of C144 and 47D11 Fab.

(a) Side and top view of a surface representation of the spike glycoprotein in pre-fusion closed (PDB ID: 6XR8), pre-fusion open (PDB ID: 6XM4) and post-fusion conformations (PDB ID: 6XRA) [58,88]. RBDs are shown in orange and glycans in pink. (b) Side and top views of the cryo-EM density maps showing C144 Fab complexed to closed pre-fusion S (top; EMD-22737) and 47D11 Fab complexed to pre-fusion S with 1 RBD up (bottom; EMD-11812) [66**,67*]. RBD in cyan and RBS in yellow. One RBS in the S protein bound with 47D11 is exposed and accessible to ACE2. Black circle in the C144-S1 complex marks the position of the loop shown in (c). Black circle in the 47D11-S complex marks the position of the loop shown in (d). (c) Enlarged view of the residues involved in the hydrophobic pocket on S protein RBD (cyan) and C144_{HC} CDRH3 loop (purple) (PDB ID: 7K90) [66**]. (d) Enlarged view of the residues involved in the hydrophobic pocket on S protein RBD (cyan) and 47D11_{HC} CDRH3 loop (purple) (PDB ID: 7AKD) [67*]. Figures were made in UCSF Chimera [87].

ACE2, and fuses the viral and cell membranes. The S trimer (Figure 3), has monomers composed of S1 and S2 domains which are cleaved by furin making the spike fusion-active. The receptor binding domain (RBD) on S1 can be either in a ‘down’ conformation hiding the receptor binding site (RBS) or in an ‘up’ conformation exposing the RBS (Figure 3a). Upon binding to ACE2, S2 undergoes conformational changes exposing the fusion peptide, and S1 dissociates leaving the post-fusion conformational state (Figure 3a) [58,59]. ACE2 glycosylation was shown to be important in its affinity for S [62]. Co-receptors include transmembrane protease serine 2 (TMPRSS2), and neuropilin binding the cleaved S [60,61].

All newly emerging SARS-CoV-2 variants of concern have amino acids changes that are being monitored in the S protein RBD (<https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/>) accessed 10th August

2021). *In vitro* evolution driven to increase S protein's affinity to ACE2 managed to recreate the emergence of the new variants of concern, Alpha, Beta and Gamma [47]. The authors showed that the ACE2 RBD domain was a highly potent inhibitor competing with membrane-bound host ACE2 for both existing strains and the newly predicted ones, confirmed by cryo-EM of existing strains [47].

One approach for inhibiting the S protein is by neutralizing antibodies (NAb), but useful epitopes must take into account the heavy surface glycosylation of S (Figure 3a). Sikora *et al.* mapped the spike protein glycans looking for unshielded epitopes using molecular simulations. This resulted in a much larger covered area than previously thought and suggests that the spike stalk is not a favorable antibody target [63]. Casalino *et al.* have done a similar mapping using all-atom molecular dynamics, finding the importance of two glycans shielding the spike RBD. The glycans play a role in changing the spike protein from the closed to the open state, as after their deletion, most spikes remained in the closed state, unable to bind to ACE2 [64*]. Hence spike production for vaccination or other functional studies on epitopes needs to be done in mammalian cells for correct glycosylation [65].


Based on the epitope location, NAb's mode of action can be competition with ACE2 for the RBS on the RBD, stabilization of the spike by binding distally to the RBS either in an 'up' or 'down' conformation of the RBD (Figure 3), and destruction of the spike by driving spike conformation to post-fusion [66**;67*;68–70]. Barnes *et al.* have tested and classified Covid19 patients isolated NAb which stabilize the spike or compete with ACE2 in three classes based on the binding site [66**]. With the emergence of new variants, a need for an antibody binding distally from the RBS arose [67*]. One of the earliest antibodies (CR3022) tested against SARS-CoV-2 was shown to potently bind the RBD at an epitope which is only exposed in the open conformation. Interestingly, using cryo-EM, no density was observed for the CR3022 FAb on the 'one up — two down' S trimer conformation which should have one epitope exposed. Instead, increasing the sample time after antibody addition yielded a homogenous population of post-fusion S proteins, suggesting that CR3022 drives S1 dissociation [69]. A similar mode of action has been observed for some Ebola antibodies, where, as with CR3022, infection is partially inhibited [69,71]. Together these findings indicate that antibodies binding to an epitope away from the RBS, stabilizing the pre-fusion closed S trimer, should be the most effective (Figure 3). Successful vaccine production is based on a similar principle, the S trimers are stabilized in a pre-fusion conformation by substitution with double proline and removal of the furin cleavage site [72]. Recent advances have been made on exploiting S proteins

stabilized in the closed conformation to produce antibodies which bind to epitopes distant from the RBS [73–76]. One such antibody is 47D11 stabilizing the S trimer in a 'one up — two down' conformation leaving one RBS accessible to ACE2 (Figure 3b down, d). It binds in a conserved hydrophobic pocket that is usually covered by glycan N343 (Figure 3d). It stabilizes the glycan in an upright position and inserts a CDRH3 loop in the pocket. Its activity was also verified against newly emerging SARS-CoV-2 variants [67*]. The C144 antibody has an epitope which is accessible in both 'up' and 'down' conformations and upon binding it spans across adjacent RBS and stabilizes the trimer in the closed conformation (Figure 3b up). It contains the same loop as 47D11, interacting with the hydrophobic pocket (Figure 3c). Its binding site still partially overlaps with the RBS (Figure 3b up) and it is not effective against all variants indicating that this approach should be explored further [66**].

Nanobodies and aptamers offer alternative, potentially cheaper, easier, faster approaches to NAb. Nanobodies are variable regions (VHH) of antibodies expressed alone. Huo *et al.* and Xiang *et al.* have shown the potential of nanobodies in inhibiting SARS-CoV-2 attachment to the host cell [77,78]. By screening through the VHH library against purified RBD Huo *et al.* discovered 2 nanobodies H11-H4 and H11-D4 with binding sites which partially overlap with the RBS. H11-H4 and H11-D4 both bind to the RBD in 'up' and 'down' conformations, stabilizing the S protein in 'one up — two down' state as shown by isothermal titration calorimetry (ITC) and X-ray crystallography [77]. Xiang *et al.* noticed this effect was increased when using multivalent nanobodies [78]. Koenig *et al.* produced multivalent nanobodies that when bound resulted in an irreversible, fully open conformation preventing S binding to the receptor. What is more, they have tested the activity in evolutionary mutants and discovered that a homotrivalent nanobody was not effective after a single mutation, however a multivalent nanobody targeting different epitopes significantly hindered the emergence of mutants [79**]. More recently, synthetic nanobodies competing with ACE2 have shown their effectiveness in hamsters [80].

Aptamers have shown potential to compete with ACE2 as nasally administered drugs [81]. For Nab, nanobodies and aptamers, the neutralization efficacy can be computationally predicted, measured through binding assays like ELISA, structural studies, pseudovirus or virus neutralization in cell culture, before going on to animal trials. Experience over the past two years has emphasized that even efficacy in ELISA does not necessarily correlate to infection inhibition in cell culture [68].

One of the central themes in inhibition of virus entry is to prevent conformational changes required to release the

Figure 4

Picornavirus' newly identified conserved site bound with inhibitor 17 bound to CVB3 and glutathione to EV-F3.

(a) Electron density map of CVB3 at 2.8 Å-resolution bound with an inhibitor (compound 17) radially colored according to the key in Å, viewed down a two-fold axis of symmetry. The position of the inhibitor is shown in orange. Two protomers are shown in light and dark grey [83*]. **(b)** Detailed view of the interprotomer position of compound 17 in CVB3, with one protomer shown in dark grey and the second protomer in light grey surface with the atomic models of VP1 (green),

VP2 (blue) and VP3 (cyan) shown in ribbon [83*]. **(c)** Close up view of compound 17 and interactions with the CVB3 site residues (PDB: 6ZCL) [83*]. **(d)** Close up view of a similar site in enterovirus-F3 as in (c) with glutathione in the position corresponding to compound 17 in CVB3 (PDB: 6T4C) [86]. (a) and (b) are adapted and reproduced with permission from Flatt et al. [83*], both under a Creative Commons Attribution 4.0 International License. (c) and (d) were made with UCSF Chimera [87].

Conclusions

Antivirals targeting vital non-structural viral proteins prevent replication or polyprotein maturation. As their targeted conserved sites are different from the host's, any antivirals targeting them should have reduced chances for off-target activity. Drug repurposing and design of broad range inhibitors are attractive approaches for finding antivirals against such targets. This has yielded a list of antivirals with partial efficacy across many targets, and sped up the search for SARS-CoV-2 inhibitors tremendously. In the case of remdesivir, it provided a quick remedy for the most critical patients in their recovery very early in the pandemic. The structural proteins, as they are exposed to the immune system, are prone to mutation. Therefore, specific inhibitors, rather than broad range inhibitors, are more common. Antivirals targeting structural proteins target the receptor attachment, membrane fusion and/or genome release, thus stopping infection at an early stage. Here, most success has been seen with antibodies, especially through the successful COVID-19 vaccine campaign targeting S. This strategy necessitates constant updating of the diagnostics, antivirals and

vaccines to ensure that variants of concern are susceptible to these proven measures. In the long run, as long as any particular virus is widespread in the population, new variants will continue to emerge, some of which will be resistant. Therefore, a combined approach that utilizes both preventative and therapeutic measures will be of benefit. To search for antivirals, virtual screening and other modeling methods have been heavily exploited. Unfortunately, most of the published modeling-oriented studies lack wet lab experimental validation. This is, in our opinion, insufficient, since any molecular modeling method alone cannot reliably predict biological activity. Commonly, molecular dynamics simulations are too short (they should be on the microsecond scale) and are carried out with too simplistic systems, for example, with many proteases, such as M^{PRO}, monomers have been used in the majority of studies, while dimers are biologically relevant. Whereas structural methods and binding affinity assays support our understanding of the mechanisms behind biological activity assays in cell culture and animal models, alone they are insufficient to infer inhibitor efficacy. One cannot forget the importance of host factors in the virus infection cycle and viral surface protein glycosylation, metastability and dissociation when designing antivirals. All of these factors may contribute to the lack of efficient antivirals so far in clinical use. The methods and models, for example, organoids used for screening, need improvement, alternative targets, especially targeting host factors that are not so prone to mutation, need to be explored more systematically, and continual monitoring of circulating virus variation should all contribute to the field in the near future.

Conflict of interest statement

Nothing declared.

Acknowledgements

The work was carried out in the Instruct Centre Finland, supported by grants from the Sigrid Jusélius Foundation (to Sarah Butcher) and the Academy of Finland (grants 336471 and 315950 to Sarah Butcher, and 336473 to Antti Poso).

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
 - of outstanding interest
- Sorokina M, Teixeira JMC, Barrera-Vilarmau S, Paschke R, Papasotiriou I, Rodrigues JPGLM, Kastritis PL: **Structural models of human ACE2 variants with SARS-CoV-2 Spike protein for structure-based drug design.** *Sci data* 2020, **7**:309.
 - Weglarz-Tomczak E, Tomczak JM, Talma M, Burda-Grabowska M, Giurg M, Brul S: **Identification of ebselein and its analogues as potent covalent inhibitors of papain-like protease from SARS-CoV-2.** *Sci Rep* 2021, **11**:3640.
 - Amin SA, Banerjee S, Singh S, Qureshi IA, Gayen S, Jha T: **First structure-activity relationship analysis of SARS-CoV-2 virus main protease (Mpro) inhibitors: an endeavor on COVID-19 drug discovery.** *Mol Divers* 2021, **25**:1827-1838 <http://dx.doi.org/10.1007/s11030-020-10166-3>.
 - Yang J, Zhang Z, Yang F, Zhang H, Wu H, Zhu F, Xue W: **Computational design and modeling of nanobodies toward SARS-CoV-2 receptor binding domain.** *Chem Biol Drug Des* 2021, **98**:1-18.
 - Tsai K-C, Lee Y-C, Tseng T-S: **Comprehensive deep mutational scanning reveals the immune-escaping hotspots of SARS-CoV-2 receptor-binding domain targeting neutralizing antibodies.** *Front Microbiol* 2021, **12**.
 - Verkhivker GM, Agajanian S, Oztas DY, Gupta G: **Comparative perturbation-based modeling of the SARS-CoV-2 spike protein binding with host receptor and neutralizing antibodies: structurally adaptable allosteric communication hotspots define spike sites targeted by global circulating mutations.** *Biochemistry* 2021, **60**:1459-1484.
 - Riva L, Yuan S, Yin X, Martin-Sancho L, Matsunaga N, Pache L, Burgstaller-Muehlbacher S, De Jesus PD, Teriete P, Hull MV et al.: **Discovery of SARS-CoV-2 antiviral drugs through large-scale compound repurposing.** *Nature* 2020, **586**:113-119.
 - Manelfi C, Gossen J, Gervasoni S, Talarico C, Albani S, Philipp BJ, Musiani F, Vistoli G, Rossetti G, Beccari AR et al.: **Combining different docking engines and consensus strategies to design and validate optimized virtual screening protocols for the SARS-CoV-2 3CL protease.** *Molecules* 2021, **26**:797.
 - Sun C, Zhang J, Wei J, Zheng X, Zhao X, Fang Z, Xu D, Yuan H, Liu Y: **Screening, simulation, and optimization design of small molecule inhibitors of the SARS-CoV-2 spike glycoprotein.** *PLoS One* 2021, **16**:e0245975.
 - Henderson JA, Verma N, Harris RC, Liu R, Shen J: **Assessment of proton-coupled conformational dynamics of SARS and MERS coronavirus papain-like proteases: Implication for designing broad-spectrum antiviral inhibitors.** *J Chem Phys* 2020, **153**:115101.
 - Ahmad S, Waheed Y, Ismail S, Najmi MH, Ansari JK: **Rational design of potent anti-COVID-19 main protease drugs: an extensive multi-spectrum in silico approach.** *J Mol Liq* 2021, **330**:115636.
 - Pavlova A, Lynch DL, Daidone I, Zanetti-Polzi L, Smith MD, Chipot C, Kneller DW, Kovalevsky A, Coates L, Golosov AA et al.: **Inhibitor binding influences the protonation states of histidines in SARS-CoV-2 main protease.** *Chem Sci* 2021, **12**:1513-1527.
 - Komatsu TS, Okimoto N, Koyama YM, Hirano Y, Morimoto G, Ohno Y, Taiji M: **Drug binding dynamics of the dimeric SARS-CoV-2 main protease, determined by molecular dynamics simulation.** *Sci Rep* 2020, **10**:16986.
 - Santana MVS, Silva-Jr FP: **De novo design and bioactivity prediction of SARS-CoV-2 main protease inhibitors using recurrent neural network-based transfer learning.** *BMC Chem* 2021, **15**:8.
 - Verma N, Henderson JA, Shen J: **Proton-coupled conformational activation of SARS coronavirus main proteases and opportunity for designing small-molecule broad-spectrum targeted covalent inhibitors.** *J Am Chem Soc* 2020, **142**:21883-21890.
 - Ke Z, Oton J, Qu K, Cortese M, Zila V, McKeane L, Nakane T, Zivanov J, Neufeldt CJ, Cerikan B et al.: **Structures and distributions of SARS-CoV-2 spike proteins on intact virions.** *Nature* 2020, **588**:498-502.
 - Huang Y, Yang C, Xu X, Xu W, Liu S: **Structural and functional properties of SARS-CoV-2 spike protein: potential antivirus drug development for COVID-19.** *Acta Pharmacol Sin* 2020, **41**:1141-1149.
 - Xiang R, Yu Z, Wang Y, Wang L, Huo S, Li Y, Liang R, Hao Q, Ying T, Gao Y et al.: **Recent advances in developing small-molecule inhibitors against SARS-CoV-2.** *Acta Pharm Sin B* 2021 <http://dx.doi.org/10.1016/j.apsb.2021.06.016>. In press.
 - McKee DL, Sternberg A, Stange U, Laufer S, Naujokat C: **Candidate drugs against SARS-CoV-2 and COVID-19.** *Pharmacol Res* 2020, **157**:104859.

20. Faheem, Kumar BK, Sekhar KVGC, Kunjiappan S, Jamalis J, Balaña-Fouce R, Tekwani BL, Sankaranarayanan M: **Druggable targets of SARS-CoV-2 and treatment opportunities for COVID-19.** *Bioorg Chem* 2020, **104**:104269.
21. Banerjee R, Perera L, Tillekeratne LMV: **Potential SARS-CoV-2 main protease inhibitors.** *Drug Discov Today* 2021, **26**:804-816.
22. Ullrich S, Nitsche C: **The SARS-CoV-2 main protease as drug target.** *Bioorg Med Chem Lett* 2020, **30**:127377.
23. Xiu S, Dick A, Ju H, Mirzaie S, Abdi F, Cocklin S, Zhan P, Liu X: **Inhibitors of SARS-CoV-2 entry: current and future opportunities.** *J Med Chem* 2020, **63**:12256-12274.
24. Zhang L, Lin D, Kusov Y, Nian Y, Ma Q, Wang J, von Brunn A, Leyssen P, Lanko K, Neyts J et al.: **α -Ketoamides as broad-spectrum inhibitors of coronavirus and enterovirus replication: structure-based design, synthesis, and activity assessment.** *J Med Chem* 2020, **63**:4562-4578.
25. Mönttinen HAM, Ravanti JJ, Poranen MM: **Structure unveils relationships between RNA virus polymerases.** *Viruses* 2021, **13**:313.
26. Boras B, Jones RM, Anson BJ, Arenson D, Aschenbrenner L, Bakowski MA, Beutler N, Binder J, Chen E, Eng H et al.: **Discovery of a novel inhibitor of coronavirus 3CL protease as a clinical candidate for the potential treatment of COVID-19.** *bioRxiv Prepr Serv Biol* 2020 <http://dx.doi.org/10.1101/2020.09.12.293498>. Submitted for publication.
27. Hillen HS: **Structure and function of SARS-CoV-2 polymerase.** *Curr Opin Virol* 2021, **48**:82-90 <http://dx.doi.org/10.1016/j.coviro.2021.03.010>.
28. Gao Y, Yan L, Huang Y, Liu F, Zhao Y, Cao L, Wang T, Sun Q, Ming Z, Zhang L et al.: **Structure of the RNA-dependent RNA polymerase from COVID-19 virus.** *Science (80-)* 2020, **368**:779-782.
29. V'kovski P, Kratzel A, Steiner S, Stalder H, Thiel V: **Coronavirus biology and replication: implications for SARS-CoV-2.** *Nat Rev Microbiol* 2021, **19**:155-170
An extensive review on SARS-CoV-2 life cycle and drug targets.
30. Jin Z, Du X, Xu Y, Deng Y, Liu M, Zhao Y, Zhang B, Li X, Zhang L, Peng C et al.: **Structure of Mpro from SARS-CoV-2 and discovery of its inhibitors.** *Nature* 2020, **582**:289-293
This research provided a starting model for designing SARS-CoV-2 main protease inhibitors based on homology with SARS-CoV.
31. Kneller DW, Galanis S, Phillips G, O'Neill HM, Coates L, Kovalevsky A: **Malleability of the SARS-CoV-2 3CL Mpro active-site cavity facilitates binding of clinical antivirals.** *Structure* 2020, **28**:1313-1320.e3.
32. Naydenova K, Muir KW, Wu L-F, Zhang Z, Coscia F, Peet MJ, Castro-Hartmann P, Qian P, Sader K, Dent K et al.: **Structure of the SARS-CoV-2 RNA-dependent RNA polymerase in the presence of favipiravir-RTP.** *Proc Natl Acad Sci USA* 2021, **118**: e2021946118.
33. Shannon A, Selisko B, Le N, Huchting J, Touret F, Piorkowski G, Fattorini V, Ferron F, Decroly E, Meier C et al.: **Favipiravir strikes the SARS-CoV-2 at its Achilles heel, the RNA polymerase.** *bioRxiv Prepr Serv Biol* 2020 <http://dx.doi.org/10.1101/2020.05.15.098731>. Submitted for publication.
34. Yin W, Mao C, Luan X, Shen D-D, Shen Q, Su H, Wang X, Zhou F, Zhao W, Gao M et al.: **Structural basis for inhibition of the RNA-dependent RNA polymerase from SARS-CoV-2 by remdesivir.** *Science (80-)* 2020, **368**:1499-1504.
35. Anand K, Palm GJ, Mesters JR, Siddell SG, Ziebuhr J, Hilgenfeld R: **Structure of coronavirus main proteinase reveals combination of a chymotrypsin fold with an extra alpha-helical domain.** *EMBO J* 2002, **21**:3213-3224.
36. Hoffman RL, Kania RS, Brothers MA, Davies JF, Ferre RA, Gajiwala KS, He M, Hogan RJ, Kozminski K, Li LY et al.: **Discovery of ketone-based covalent inhibitors of coronavirus 3CL proteases for the potential therapeutic treatment of COVID-19.** *J Med Chem* 2020, **63**:12725-12747.
37. Sacco MD, Ma C, Lagarias P, Gao A, Townsend JA, Meng X, Dube P, Zhang X, Hu Y, Kitamura N et al.: **Structure and inhibition of the SARS-CoV-2 main protease reveal strategy for developing dual inhibitors against Mpro and cathepsin L.** *Sci Adv* 2020, **6**.
38. Dai W, Zhang B, Jiang X-M, Su H, Li J, Zhao Y, Xie X, Jin Z, Peng J, Liu F et al.: **Structure-based design of antiviral drug candidates targeting the SARS-CoV-2 main protease.** *Science* 2020, **368**:1331-1335.
39. Mengist HM, Dilnessa T, Jin T: **Structural basis of potential inhibitors targeting SARS-CoV-2 main protease.** *Front Chem* 2021, **9**.
40. Mengist HM, Mekonnen D, Mohammed A, Shi R, Jin T: **Potency, safety, and pharmacokinetic profiles of potential inhibitors targeting SARS-CoV-2 main protease.** *Front Pharmacol* 2021, **11**.
41. Zhang L, Lin D, Sun X, Curth U, Drosten C, Sauerhering L, Becker S, Rox K, Hilgenfeld R: **Crystal structure of SARS-CoV-2 main protease provides a basis for design of improved α -ketoamide inhibitors.** *Science (80-)* 2020, **368**:409-412.
42. Lockbaum GJ, Reyes AC, Lee JM, Tilawala R, Nalivaika EA, Ali A, Kurt Yilmaz N, Thompson PR, Schiffer CA: **Crystal structure of SARS-CoV-2 main protease in complex with the non-covalent inhibitor ML188.** *Viruses* 2021, **13**:174.
43. Tian L, Qiang T, Liang C, Ren X, Jia M, Zhang J, Li J, Wan M, YuWen X, Li H et al.: **RNA-dependent RNA polymerase (RdRp) inhibitors: the current landscape and repurposing for the COVID-19 pandemic.** *Eur J Med Chem* 2021, **213**:113201.
44. Yin W, Luan X, Li Z, Zhou Z, Wang Q, Gao M, Wang X, Zhou F, Shi J, You E et al.: **Structural basis for inhibition of the SARS-CoV-2 RNA polymerase by suramin.** *Nat Struct Mol Biol* 2021, **28**:319-325.
45. Picarazzi F, Vicenti I, Saladini F, Zazzi M, Mori M: **Targeting the RdRp of emerging RNA viruses: the structure-based drug design challenge.** *Molecules* 2020, **25**:5695.
46. Curreli F, Victor SMB, Ahmed S, Drelich A, Tong X, Tseng C-TK, Hillyer CD, Debnath AK: **Stapled peptides based on human angiotensin-converting enzyme 2 (ACE2) potently inhibit SARS-CoV-2 infection in vitro.** *mBio* 2020, **11**.
47. Zahradník J, Marciano S, Shemesh M et al.: **SARS-CoV-2 variant prediction and antiviral drug design are enabled by RBD in vitro evolution.** *Nat Microbiol* 2021, **6**:1188-1198 <https://doi.org/10.1038/s41564-021-00954-4>.
48. Oliveira MA, Zhao R, Lee W-M, Kremer MJ, Minor I, Rueckert RR, Diana GD, Pevsar DC, Dutko FJ, McKinlay MA et al.: **The structure of human rhinovirus 16.** *Structure* 1993, **1**:51-68.
49. Sherman MB, Smith HQ, Smith TJ: **The dynamic life of virus capsids.** *Viruses* 2020, **12**:618.
50. Linden L, Wolthers K, van Kuppeveld F: **Replication and inhibitors of enteroviruses and parechoviruses.** *Viruses* 2015, **7**:4529-4562.
51. Baker JD, Uhrich RL, Kraemer GC, Love JE, Kraemer BC: **A drug repurposing screen identifies hepatitis C antivirals as inhibitors of the SARS-CoV2 main protease.** *PLoS One* 2021, **16**:e0245962.
52. Anand K: **Coronavirus main proteinase (3CLpro) Structure: basis for design of anti-SARS drugs.** *Science (80-)* 2003, **300**:1763-1767.
53. Yang H, Xie W, Xue X, Yang K, Ma J, Liang W, Zhao Q, Zhou Z, Pei D, Ziebuhr J et al.: **Design of wide-spectrum inhibitors targeting coronavirus main proteases.** *PLoS Biol* 2005, **3**:e324.
54. Weng YL, Naik SR, Dingelstad N, Lugo MR, Kalyaanamoorthy S, Ganesh A: **Molecular dynamics and in silico mutagenesis on the reversible inhibitor-bound SARS-CoV-2 main protease complexes reveal the role of lateral pocket in enhancing the ligand affinity.** *Sci Rep* 2021, **11**:7429.
55. Arora R, Liew CW, Soh TS, Otoo DA, Seh CC, Yue K, Nilar S, Wang G, Yokokawa F, Noble CG et al.: **Two RNA tunnel inhibitors**

- bind in highly conserved sites in dengue virus NS5 polymerase: structural and functional studies.** *J Virol* 2020, **94**.
56. Yang J, Jing X, Yi W, Li X-D, Yao C, Zhang B, Zheng Z, Wang H, Gong P: **Crystal structure of a tick-borne flavivirus RNA-dependent RNA polymerase suggests a host adaptation hotspot in RNA viruses.** *Nucleic Acids Res* 2021, **49**:1567-1580.
57. Gejji V, Svoboda P, Stefanik M, Wang H, Salat J, Eyer L, Ruzek D, Fernando S: **An RNA-dependent RNA polymerase inhibitor for tick-borne encephalitis virus.** *Virology* 2020, **546**:13-19.
58. Cai Y, Zhang J, Xiao T, Peng H, Sterling SM, Walsh RM, Rawson S, Rits-Volloch S, Chen B: **Distinct conformational states of SARS-CoV-2 spike protein.** *Science (80-)* 2020, **369**:1586-1592.
59. Walls AC, Park Y-J, Tortorici MA, Wall A, McGuire AT, Veesler D: **Structure, function, and antigenicity of the SARS-CoV-2 spike glycoprotein.** *Cell* 2020, **181**:281-292.e6.
60. Cantuti-Castelvetro L, Ojha R, Pedro LD, Djannatian M, Franz J, Kuivanen S, van der Meer F, Kallio K, Kaya T, Anastasina M et al.: **Neuropilin-1 facilitates SARS-CoV-2 cell entry and infectivity.** *Science* 2020, **370**:856-860.
61. Daly JL, Simonetti B, Klein K, Chen K-E, Williamson MK, Antón-Plágaro C, Shoemark DK, Simón-Gracia L, Bauer M, Hollandi R et al.: **Neuropilin-1 is a host factor for SARS-CoV-2 infection.** *Science* 2020, **370**:861-865.
62. Mehdipour AR, Hummer G: **Dual nature of human ACE2 glycosylation in binding to SARS-CoV-2 spike.** *Proc Natl Acad Sci U S A* 2021, **118**:e2100425118.
63. Sikora M, von Bülow S, Blanc FEC, Gecht M, Covino R, Hummer G: **Computational epitope map of SARS-CoV-2 spike protein.** *PLoS Comput Biol* 2021, **17**:e1008790.
64. Casalino L, Gaieb Z, Goldsmith JA, Hjorth CK, Dommer AC, Harbison AM, Fogarty CA, Barros EP, Taylor BC, McLellan JS et al.: **Beyond shielding: the roles of glycans in the SARS-CoV-2 spike protein.** *ACS Cent Sci* 2020, **6**:1722-1734
- This research identified two important glycans for the SARS-CoV-2 spike protein function. These could be potential new targets and important considerations in antibody and nanobody design.
65. Brun J, Vasiljevic S, Gangadharan B, Hensen M, Chandran AV, Hill ML, Kiappes JL, Dwek RA, Alonzi DS, Struwe WB et al.: **Assessing antigen structural integrity through glycosylation analysis of the SARS-CoV-2 viral spike.** *ACS Cent Sci* 2021, **7**:586-593.
66. Barnes CO, Jette CA, Abernathy ME, Dam K-MA, Esswein SR, Gristick HB, Malyutin AG, Sharaf NG, Huey-Tubman KE, Lee YE et al.: **SARS-CoV-2 neutralizing antibody structures inform therapeutic strategies.** *Nature* 2020, **588**:682-687
- Authors show different antibodies' modes of action and binding sites on the spike and classify them making the further research on stabilizing S protein by antibodies more direct.
67. Fedry J, Hurdiss DL, Wang C, Li W, Obal G, Drulyte I, Du W, Howes SC, van Kuppeveld FJM, Förster F et al.: **Structural insights into the cross-neutralization of SARS-CoV and SARS-CoV-2 by the human monoclonal antibody 47D.** *Sci Adv* 2021, **7**:eabf5632
- Authors show the binding of a cross-neutralizing antibody to SARS-CoV and SARS-CoV-2 S protein and elucidate further on the differences in the S protein across Coronavirus genus and among SARS-CoV-2 variants.
68. Asarnow D, Wang B, Lee W-H, Hu Y, Huang C-W, Faust B, Ng PML, Ngoh EZX, Bohn M, Bulkley D et al.: **Structural insight into SARS-CoV-2 neutralizing antibodies and modulation of syncytia.** *Cell* 2021, **184**:3192-3204.e16.
69. Huo J, Zhao Y, Ren J, Zhou D, Duyvesteyn HME, Ginn HM, Carrique L, Malinauskas T, Ruza RR, Shah PNM et al.: **Neutralization of SARS-CoV-2 by Destruction of the Prefusion Spike.** *Cell Host Microbe* 2020, **28**:497.
70. Wu Y, Wang F, Shen C, Peng W, Li D, Zhao C, Li Z, Li S, Bi Y, Yang Y et al.: **A noncompeting pair of human neutralizing antibodies block COVID-19 virus binding to its receptor ACE2.** *Science* 2020, **368**:1274-1278.
71. Rijal P, Elias SC, Machado SR, Xiao J, Schimanski L, O'Dowd V, Baker T, Barry E, Mendelsohn SC, Cherry CJ et al.: **Therapeutic monoclonal antibodies for Ebola virus infection derived from vaccinated humans.** *Cell Rep* 2019, **27**:172-186.e7.
72. Amanat F, Strohmeier S, Rathnasinghe R, Schotsaert M, Coughlan L, Garcia-Sastre A, Krammer F: **Introduction of two prolines and removal of the polybasic cleavage site lead to higher efficacy of a recombinant spike-based SARS-CoV-2 vaccine in the mouse model.** *mBio* 2021, **12**.
73. Carnell GW, Ciazynska KA, Wells DA, Xiong X, Aguinam ET, McLaughlin SH, Mallery D, Ebrahimi S, Ceron-Gutierrez L, Asbach B et al.: **SARS-CoV-2 spike protein stabilized in the closed state induces potent neutralizing responses.** *J Virol* 2021, **95**.
74. Xiong X, Qu K, Ciazynska KA, Hosmillo M, Carter AP, Ebrahimi S, Ke Z, Scheres SHW, Bergamaschi L, Grice GL et al.: **A thermostable, closed SARS-CoV-2 spike protein trimer.** *Nat Struct Mol Biol* 2020, **27**:934-941.
75. McCallum M, Walls AC, Bowen JE, Corti D, Veesler D: **Structure-guided covalent stabilization of coronavirus spike glycoprotein trimers in the closed conformation.** *Nat Struct Mol Biol* 2020, **27**:942-949.
76. Henderson R, Edwards RJ, Mansouri K, Janowska K, Stalls V, Gobeil SMC, Kopp M, Li D, Parks R, Hsu AL et al.: **Controlling the SARS-CoV-2 spike glycoprotein conformation.** *Nat Struct Mol Biol* 2020, **27**:925-933.
77. Huo J, Le Bas A, Ruza RR, Duyvesteyn HME, Mikolajek H, Malinauskas T, Tan TK, Rijal P, Dumoux M, Ward PN et al.: **Neutralizing nanobodies bind SARS-CoV-2 spike RBD and block interaction with ACE2.** *Nat Struct Mol Biol* 2020, **27**:846-854.
78. Xiang Y, Nambulli S, Xiao Z, Liu H, Sang Z, Duprex WP, Schneidman-Duhovny D, Zhang C, Shi Y: **Versatile and multivalent nanobodies efficiently neutralize SARS-CoV-2.** *Science* 2020, **370**:1479-1484.
79. Koenig P-A, Das H, Liu H, Kümmeler BM, Gohr FN, Jenster L-M, Schiffelers LDJ, Tesfamariam YM, Uchima M, Wuerth JD et al.: **Structure-guided multivalent nanobodies block SARS-CoV-2 infection and suppress mutational escape.** *Science* 2021, **371**
- Authors show the potential of multivalent nanobodies in tackling the inhibition of new emerging strains of SARS-CoV-2.
80. Li T, Cai H, Yao H, Zhou B, Zhang N, van Vliessingen MF, Kuiken T, Han W, GeurtsvanKessel CH, Gong Y et al.: **A synthetic nanobody targeting RBD protects hamsters from SARS-CoV-2 infection.** *Nat Commun* 2021, **12**:4635.
81. Sun M, Liu S, Wei X, Wan S, Huang M, Song T, Lu Y, Weng X, Lin Z, Chen H et al.: **Aptamer blocking strategy inhibits SARS-CoV-2 virus infection.** *Angew Chemie Int Ed* 2021, **60**:10266-10272.
82. Abdelnabi R, Geraets JA, Ma Y, Mirabelli C, Flatt JW, Domanska A, Delang L, Jochmans D, Kumar TA, Jayaprakash V et al.: **A novel druggable interprotomer pocket in the capsid of rhino- and enteroviruses.** *PLoS Biol* 2019, **17**:e3000281.
83. Flatt JW, Domanska A, Seppälä AL, Butcher SJ: **Identification of a conserved virion-stabilizing network inside the interprotomer pocket of enteroviruses.** *Commun Biol* 2021, **4**:250
- The authors show the inhibition of coxsackievirus B3 and B4 with small molecules binding to a conserved, interprotomer, druggable capsid pocket which could be used in drug development against enteroviruses for which there are no antivirals currently.
84. Ruokolainen V, Domanska A, Laajala M, Pelliccia M, Butcher SJ, Marjomäki V: **Extracellular albumin and endosomal ions prime enterovirus particles for uncoating that can be prevented by fatty acid saturation.** *J Virol* 2019, **93**.
85. Shah PNM, Filman DJ, Karunatilaka KS, Hesketh EL, Groppelli E, Strauss M, Hogle JM: **Cryo-EM structures reveal two distinct conformational states in a picornavirus cell entry intermediate.** *PLoS Pathog* 2020, **16**:e1008920.
86. Duyvesteyn HME, Ren J, Walter TS, Fry EE, Stuart DI: **Glutathione facilitates enterovirus assembly by binding at a druggable pocket.** *Commun Biol* 2020, **3**:9.

87. Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE: **UCSF Chimera—a visualization system for exploratory research and analysis.** *J Comput Chem* 2004, **25**:1605-1612.
88. Zhou Tongqing, Tsybovsky Yaroslav, Olia Adam S, Gorman Jason, Rapp Micah, Cerutti Gabriele, Chuang Gwo-Yu, Katsamba Phinikoula S, Nazzari Alexandra, Sampson Jared M et al.: **Cryo-EM structures delineate a pH-dependent switch that mediates endosomal positioning of SARS-CoV-2 spike receptor-binding domains.** *Cell Host & Microbe* 2020, **28**:867-879 <http://dx.doi.org/10.1016/j.chom.2020.11.004>.